

Multi-sensor Improved Sea Surface Temperature (MISST) for GODAE

Lead PI : Chelle L. Gentemann
438 First St, Suite 200; Santa Rosa, CA 95401-5288
Phone: (707) 545-2904x14 FAX: (707) 545-2906 E-mail: gentemann@remss.com

CO-PI: Gary A. Wick
NOAA/ETL R/ET6, 325 Broadway, Boulder, CO 80305
Phone: (303) 497-6322 FAX: (303) 497-6181 E-mail: gary.a.wick@noaa.gov

CO-PI: James Cummings
Oceanography Division, Code 7320, Naval Research Laboratory, Monterey, CA 93943
Phone: (831) 656-5021 FAX: (831) 656-4769 E-mail: cummings@nrlmry.navy.mil

CO-PI: James Yoe
NOAA/NESDIS, E/RA2 Room 808 WWB, 5200 Auth Road, Camp Springs, MD 20746-4304
Phone: (301) 763-8172x186 FAX: (301) 763-8149 E-mail: James.G.Yoe@noaa.gov

Award Number: *NNG04GM56G*
<http://www.misst.org>

LONG-TERM GOALS

The Multi-sensor Improved Sea Surface Temperatures (MISST) for the Global Ocean Data Assimilation Experiment (GODAE) project intends to produce an improved, high-resolution, global, near-real-time (NRT), sea surface temperature analysis through the combination of satellite observations from complementary infrared (IR) and microwave (MW) sensors and to then demonstrate the impact of these improved sea surface temperatures (SSTs) on operational ocean models, numerical weather prediction, and tropical cyclone intensity forecasting.

SST is one of the most important variables related to the global ocean-atmosphere system. It is a key indicator for climate change and is widely applied to studies of upper ocean processes, to air-sea heat exchange, and as a boundary condition for numerical weather prediction. The importance of SST to accurate weather forecasting of both severe events and daily weather has been increasingly recognized over the past several years. Despite the importance and wide usage of operational SST analyses, significant weaknesses remain in the existing operational products.

The improved sensors on the Terra, Aqua, and EnviSAT-1 satellites, in conjunction with previously existing sensors on several other US Navy, NASA, and NOAA satellites, provide the opportunity for notable advances in SST measurement. In addition to more frequent coverage for increased temporal resolution, these sensors permit the combination of highly complementary IR and MW retrievals. While clouds, aerosols, and atmospheric water vapor affect IR retrievals, these phenomena have little impact on MW retrievals. Characteristically, IR SST provides high spatial resolution (~1 km at nadir) but poorer coverage with the presence of clouds. Although having a reduced resolution (~25 km grid), MW SST provide >90% coverage of the global ocean each day. These factors have motivated interest in the development of merged IR and MW SST products to leverage the positive characteristics of each

sensor type. Merging multiple SST sensors into a single analysis will result in enhanced reliability, availability, and accuracy.

This project has two distinct goals: (1) producing an improved sea surface temperature (SST) product through the combination of observations from complementary infrared (IR) and microwave (MW) sensors, and (2) demonstrating the impact of improved multi-sensor SST products on operational ocean models, numerical weather prediction, and tropical cyclone intensity forecasting. Close collaboration and the international coordinated exchange of SST products with error statistics with operational agencies will optimize utility of these new data streams by US and international operational agencies. Innovative techniques to blend these complementary data will be applied in operational frameworks at NOAA and Navy. This project will make a direct US contribution to the Global Ocean Data Assimilation Experiment (GODAE) by working within the GODAE High-Resolution SST Pilot Project (GHRSSST-PP), initiated by the international GODAE steering team, to coordinate the production of a new generation high-resolution SST. By contributing to the GHRSSST-PP this team will minimize duplication of efforts, harmonize research and development activities, and maximize data access.

This effort will ensure that US scientists and operational activities remain at the forefront of the international ocean and weather forecasting activities and are provided with state-of-the-art SST data products and analyses.

OBJECTIVES

To produce multi-sensor improved SSTs and successfully assess the impact of these products, five clear project objectives have been identified:

- 1) Computation of sensor-specific observational error characteristics is required for optimal application and data fusion techniques.
- 2) Parameterization of IR and MW retrieval differences, with consideration of diurnal warming of the ocean surface and cool-skin effects at the air-sea interface is required for multi-sensor blending and production of both skin and bulk analyses.
- 3) Production and dissemination of Level 2 Processed (L2P) sensor-specific SST products with associated retrieval bias error, standard deviation (STD), and diurnal warming estimates to the application user community.
- 4) Production and dissemination of improved multi-sensor high-resolution skin and bulk SST analyses to demonstrate and optimize utility in operational applications.
- 5) Targeted impact assessment of the SST analyses on hurricane intensity forecasting, numerical data assimilation by ocean models (both national and within GODAE), numerical weather prediction, and operational ocean forecast models.

APPROACH AND WORK PLAN

Production of a multi-sensor, improved SST product requires detailed, consistent processing of all input data and characterization of retrieval errors and differences in addition to development of fusion techniques. Much of the methodology to be applied is selected for consistency with the GHRSSST-PP Data processing Specifications (GDS), which is being designed to produce SST data products that satisfy the requirements of existing operational ocean forecast and prediction systems.

This project will also provide an assessment of the operational impact of improvements by the enhanced sampling and error characterization of the IR and MW sensors in the areas of NWP and ocean modeling. Targeted applications include Navy fleet operations, naval and civilian NWP, operational oceanography, and climate monitoring and forecasting. Each of these areas is of national importance and has corresponding national programs. For each of these applications, it is anticipated that this project will provide significant enhancements to the quality and availability of data. Through affiliation with the GHRSS-PP, the products will also be directly utilized by the international GODAE modeling communities. This product sharing will be achieved through the partnerships and close connections between the data provider and user communities.

The MISST project has a broad partnership of scientists from academia, government, and private industry, including **Remote Sensing Systems** (C. Gentemann), **NOAA** (G. Wick, J. Cione, K. Casey, J. Yoe, M. DeMaria, R. Reynolds), **NRL** (J. Cummings, C. Barron, N. Baker, J. Goerss), **NAVOCEANO** (D. May), **NASA JPL PODAAC** (J. Vasquez), **U. Maryland** (A. Harris), **U. Edinburgh** (C. Merchant), **U. Miami** (P. Minnett, B. Evans, E. Chassignet), **U. Colorado** (B. Emery, S. Castro), **WHOI** (B. Ward), and the **International GHRSS-PP Project Office** (C. Donlon). The project has identified specific tasks: data provision (C. Gentemann, B. Evans, D. May, J. Yoe), determining sensor errors (G. Wick, C. Gentemann, A. Harris, P. Minnett, B. Evans, S. Castro, B. Emery, D. May), modeling diurnal warming and skin layer effects (G. Wick, C. Gentemann, S. Castro, B. Emery, B. Ward), production and distribution of data analyses (J. Cummings, J. Yoe, J. Vasquez), and performing impact studies (J. Yoe, J. Cione, J. Goerss, N. Baker, E. Chassignet, C. Barron).

Additionally, this NOPP project is closely partnered with two other NOPP projects: “U.S. GODAE: Global Ocean Prediction with the HYbrid Coordinate Ocean Model (HYCOM)” and “POSITIV: Prototype Operational System – ISAR – Temperature Instrumentation for the VOS fleet”. The HYCOM Consortium will evaluate the impact of the improved SSTs on its ocean forecast system and the POSITIV ocean temperatures will be utilized in NRT for error characterization of satellite SST retrievals.

During the third year of the project (June 2005 – June 2006) we plan to (1) continue the NRT *in situ* satellite database, (2) continue development of error characteristics for MODIS, AVHRR, GOES, TMI, and AMSR-E SSTs (3) produce L2P for AVHRR, TMI, and AMSR-E, (4) update the diurnal and skin layer modules, (5) continue to produce blended IR/MW SST, (6) continue validation of blended SSTs, (7) begin impact studies.

WORK COMPLETED

(1) Collocated satellite *in situ* database are established and continually updated for AVHRR, MODIS, AMSR-E, and TMI, available in NRT from NAVOCEANO, U. Miami, and RSS. (2) Updated reports on the error characteristics for MODIS, AVHRR, GOES, TMI, and AMSR-E SSTs. (3) Orbital and/or gridded L2P and associated support files are available in NRT for AVHRR, GOES, TMI, and AMSR-E. (4) Completed initial diurnal warming model evaluations and sensitivity studies, updates to the diurnal and skin layer modules are completed, but research continues in this area. (5) Blended IR/MW SST products from RSS (MODIS, TMI, and AMSR-E), NRL (AVHRR and AMSR-E), NOAA (AVHRR and GOES). (6) Validation of blended SSTs is underway, (7) NAVOCEANO’s implementation of NCODA has been updated to assimilate additional data (AVHRR GAC, AVHRR LAC, GOES, AMSR-E, AATSR, and MSG SSTs). (8) Impact studies are on-going: NRL – evaluate impact of MISST SSTs on Navy Operational Global Atmospheric Prediction System (NOGAPS)

hurricane forecasts, NRL – evaluate impact of MISST SSTs on ocean models, specifically the synthetic profiles of temperature and salinity used to estimate the sound speed field for underwater acoustic applications, NOAA - evaluate impact of MISST SSTs on the Statistical Hurricane Intensity Prediction System (SHIPS), NOAA – evaluate impact of MISST SSTs on NCEP operational weather forecasts and ocean prediction.

RESULTS

SST error estimates

Significant progress has been made towards a better understanding of the errors in MW and IR SST retrievals and methods to estimate these errors in NRT. Better estimates of sensor errors are necessary for multi-sensor data fusion because the accuracy of the commonly used statistical method depends on the accuracy of our understanding and modeling of biases between sensors. Multiple initial proposed methodologies for the computation of error estimates for the AVHRR were inter-compared using extended collocations with drifting and moored buoys. The effectiveness of the formulations was objectively assessed through application to direct differencing of IR and MW data. The work yielded a multi-tiered look-up table for the errors where different optimal combinations of inputs are used depending on data availability. Different methodologies are recommended based on whether the application is to real-time processing or reanalysis.

U. Miami/RSMAS has developed error arrays, analogous to those previously produced for AVHRR, for MODIS AQUA and TERRA. These are based on MAERI radiometric and buoy in situ observations coupled to the complete mission lifetime for the two MODIS sensors. Error statistics computed from these error 'hypercubes' will be used to compute expected bias and standard deviation needed for MODIS AQUA and TERRA SST L2P datasets.

TMI and AMSR-E error statistics were computed using both *in situ* data and the Reynolds optimum interpolated SST. The errors are calculated based on NRT buoy collocations and look-up tables with error determined as a function of SST, wind speed, and distance to land.

NOAA research has determined that estimation of the bias caused by aerosols in infrared SST retrievals can most optimally be calculated using a fast-forward radiative transfer model (RTM) and inclusion of the full aerosol field in the state vector. Based on NOAA research results, estimates of aerosol optical depth has been added to the NAVOCEANO data stream. Wavelength-dependent aerosol optical depth information is now being generated by the Navy Aerosol Analysis and Prediction System and full 3-D fields are being made available to test the hypothesis.

NAVOCEANO has developed a methodology to add information on retrieval error to the US NAVY operational data stream. This method adds quantitative estimates of reliability to every MCSST sample operationally generated at NAVOCEANO. The current scheme appears to be robust and low-maintenance, both attributes being major requirements in an operational environment. Retrieval errors are calculated for N-16, N-17, and N-18 AVHRR GAC and LAC data.

Evaluation of the error estimates from the MISST data providers is being performed at FNMOC. Results showed that the AVHRR GAC error estimates are consistent with the innovations, but that the AMSR-E error estimates from the data provider are too low. A representation error computed from the gradient of the background field times the ratio of the spatial averaging properties of the SST retrieval to the grid resolution was added to the error estimates for both the AVHRR GAC and AMSR-E MW

retrievals. The diagnostic for this run was consistent with the innovation time series. Further work on the sensitivity of the J_{\min} diagnostic to the MISST error estimates from the data providers is on going.

Datasets

Several new SST datasets, in a common format, are available through this project. NAVOCEANO is producing N-18 AVHRR GAC and LAC L2P files. RSS is producing both gridded and orbital TMI and AMSR-E L2P files. NOAA is producing GOES L2P files. This new SST data format has SST observations with ancillary fields that are necessary for optimal use of the data within SST analysis systems. They also provide a US contribution to the international data sharing of SST data in the L2P format.

Diurnal Warming and Skin Models

Comparisons of simulations from full diurnal warming models with detailed temperature profile measurements from the SkinDeEP instrument identified the best combination of model physics for use in evaluation of simplified parameterizations and quantified expected errors in estimates of diurnal warming. Comparison of diurnal models to *in situ* data has resulted in improvements in modeling capabilities.

NOAA has developed a new skin model was developed using a large database of skin observations. This model is used to estimate the skin effect missing from MW SSTs, and is included in the L2P data produced by RSS.




Figure 1. FNMOC high resolution (9 km) SST on February 14, 2006.

Global SST publicly distributed (all data available at www.misst.org)

Several global blended MW and MW/IR SSTs are available from RSS in NRT. Three blended products are available at daily 25-km resolution. The MW/IR daily 9-km global SST utilizes observations from TMI, AMSR-E, and MODIS. Initial validation results show that the cloud contamination in MODIS remains an issue and regional differences between the IR and MW data need to be better parameterized, but the blending of MW and IR SSTs is leading to a more accurate, higher resolution SST than previously possible.

A Global SST and sea ice analysis is available from FNMOC (Figure 1). The analysis is executed on a 9-km mid-latitude (12-km at the equator) resolution grid and is updated every six hours. SST

observations used in the analysis include GAC retrievals from AVHRR (NOAA-16, 17, 18) and in situ data from ships and fixed and drifting buoys.

Global SST under-development

A global MW/IR SST and sea ice analysis is under development at FNMOC. The 9-km operational analysis will be used as the baseline Navy system for further development. Initial efforts toward a blended MW/IR SST analysis product indicate that additional quality control is required on the AMSR-E data by comparison against the AVHRR/in situ SST analysis product valid at the time of the microwave SST retrieval. Although the AMSR-E rejection rates were very low (~0.3% over a three month period from July – August, 2005), some rejected retrievals had anomalies greater than 4°C from the background field. The source of these large AMSR-E anomalies is under investigation.

A global blended AVHRR and GOES SST is under development at NOAA. Internal validation of the product is underway.

IMPACT AND APPLICATIONS

National Security

SST is routinely used both directly in Naval fleet operations and as an input to weather forecast models used to support Naval operations. The improved SST products and better understanding of the associated errors resulting from this project will provide a more accurate description of environmental conditions enabling better planning of operations. A key aspect of this project is directly evaluating the impact of the improved SSTs on Naval applications. SSTs are also a key parameter for identifying the location and strengths of thermal fronts and eddies, information crucial to assessing the acoustic environment for submarine and antisubmarine operations, as well as for Homeland Security considerations of coastal currents and eddies for public health and safety in the advent of deliberate dumping and dispersion of hazardous material.

Economic Development

SST data is a significant consideration for planning and conducting commercial fishing operations, as well as fisheries management and monitoring efforts. Likewise, SST data is relevant to marine protected species monitoring and de-conflicting protection efforts from commercial fishing.

Quality of Life

The potential for producing more accurate SST products has important application to areas including environmental monitoring and weather forecasting. More accurate knowledge of the SST can lead to improved understanding of coral health, better forecasting of routine and severe weather events, improved recreational fishing, and increased ability to monitor climate change. Improved understanding in these areas will lead to a more informed public and better decision-making. The specific focus on tropical cyclone intensity forecasting will potentially impact warning and evacuation decisions.

Science Education and Communication

The NASA Earth Observatory (www.earthobservatory.nasa.gov) provides an online magazine on NASA Earth Sciences that includes feature articles, daily news and images, breaking news Earth Sciences events, as well as providing global sets for online browsing. MISST SST analyses are

utilized daily to provide visuals for a variety of media updates and alerts, the most common of which are hurricane-related activities. NASA Earth Observatory provides support to a number of museum projects by generating automated near-real time data sets. Among these, the Forces of Change exhibit at the National (Smithsonian) Museum of Natural History is prototypical. We provided four near-real time data sets sized and co-registered to their system requirements, of which one is the MISST SST analysis. The Forces of Change was a temporary exhibit, but a data kiosk at NMNH continues to draw on these data and visualizations, and a number of other museums with parallel interests (e.g. the Science Museum of Brooklyn) routinely pick up this near-real time data, as well as it being quite useful for periodic requests from NASA/GSFC Public Affairs Office (PAO), staff scientists wanting to talk about ENSO events with reporters, etc. The MISST data has also been appearing in such flagship NASA internal productions such as NASA's Hurricane Watch (http://www.nasa.gov/mission_pages/hurricanes/main/index.html), of which one of the most intriguing visualizations uses MISST SST, GOES clouds, and recorded storm tracks to show the 2005 hurricane season: 27 storms Arlene to Zeta (<http://learners.gsfc.nasa.gov/mediaviewer/27storms/>).

TRANSITIONS

National Security

Through direct project partnership with US Navy efforts, the improved SST products and methodologies will be directly integrated into Naval SST products and numerical weather forecasting procedures both in use and under evaluation. As one example, improved error characterization will be incorporated in the NCODA model. To accomplish the goal of determining the impact of the SST improvements in Naval applications, transitioning results to the Naval partners is a central focus of this project.

Economic Development

Satellite IR SST data are already in use by the National Marine Fisheries Service. Improved coverage in persistently cloudy regions will facilitate protected species and fisheries management efforts. The merged IR-MW SST product will be provided when available via the NOAA CoastWatch program.

Quality of Life

Key impact assessments are planned in the areas of numerical weather prediction and tropical cyclone intensity forecasting through the activities of project partners. New SST retrievals and improved error estimates are to be integrated into existing forecast models to determine their impact. Additionally, the merged products will be provided to NOAA's National Weather Service (NWS) National Centers for Environmental Prediction (NCEP)'s Ocean Prediction Center to support ocean forecasting of winds and waves, as well as thermal conditions. Through involvement with the international GHRSSST-PP, the resulting products will be further available for incorporation by a diverse, interested user group.

RELATED PROJECTS

“U.S. GODAE: Global Ocean Prediction with the HYbrid Coordinate Ocean Model (HYCOM)”: <http://hycom.rsmas.miami.edu/>

“POSITIV: Prototype Operational System – ISAR – Temperature Instrumentation for the VOS fleet”